

Table 4. Evaluation of water components of various cropping sequences

Cropping sequence	Irrigation water applied (cm)	Profile water use (cm)	Rainfall (cm)	Water expense (cm)	Water productivity (kg m ⁻³)
Cotton-wheat	67.5	-10.88	43.56	100.18	0.327
Guar-wheat	45.0	7.59	32.19	84.78	0.358
Moong-wheat	52.5	-3.97	30.81	79.34	0.273
Cotton-barley	60	8.94	43.56	94.62	0.286
Guar- barley	37.5	9.51	32.19	79.20	0.323
Moong-barley	45.0	-1.8	30.81	74.01	0.227
Cotton- <i>raya</i>	60.0	-8.39	43.56	95.17	0.271
Guar- <i>raya</i>	37.5	8.84	32.19	78.53	0.333
Moong- <i>raya</i>	45.0	-1.6	30.81	74.21	0.242

Table 5. Effect of various cropping sequences on soil properties (0-15 cm) after completion of the experiment

Crop Sequence	pH (1:2)	EC (1:2) (dSm ⁻¹)	OC (%)	Av. N (kg ha ⁻¹)	AV. P (kg ha ⁻¹)	Av. K (kg ha ⁻¹)
Cotton-wheat	8.49	0.213	0.30	107	16.8	312
Guar-wheat	8.46	0.173	0.37	121	18.9	332
Moong-wheat	8.40	0.167	0.33	117	18.0	330
Cotton-barley	8.48	0.187	0.30	111	16.6	315
Guar-barley	8.43	0.147	0.38	123	18.4	339
Moong-barley	8.36	0.148	0.35	118	17.8	337
Cotton- <i>raya</i>	8.45	0.206	0.29	108	16.3	312
Guar- <i>raya</i>	8.34	0.149	0.35	118	18.5	325
Moong- <i>raya</i>	8.40	0.148	0.33	117	18.2	320
SEm ±	0.81	0.006	0.012	1.12	0.95	2.1
CD (5%)	-	-	0.06	5.2	-	7.7

Table 6. Economics of the various cropping sequences in comparison with existing sequence

Cropping sequence	Variable cost (Rs ha ⁻¹)	Gross returns (Rs ha ⁻¹)	Net returns (Rs ha ⁻¹)	Increase/decrease over cotton-wheat sequence (Rs ha ⁻¹)	Benefit : cost ratio
Cotton-wheat	70901	97192	26291	-	1.37
Guar-wheat	37772	90063	52291	+26000	2.36
Moong-wheat	47777	64266	16489	-9802	1.34
Cotton-barley	62427	80298	17871	-8420	1.29
Guar- barley	29298	75884	46586	+20295	2.56
Moong-barley	39303	49848	10545	-15746	1.27
Cotton- <i>raya</i>	62819	76505	13687	-12604	1.22
Guar- <i>raya</i>	29690	77589	47900	+21609	2.59
Moong- <i>raya</i>	39695	53299	13604	-12687	1.34

sequence. The potassium content decreased in cotton based crop sequence (312 kg ha⁻¹) from initial status (326 kg ha⁻¹) but, slight increase in available K was noticed in *guar*-barley (339 kg ha⁻¹), *moong*-barley (337 kg ha⁻¹), *guar*-wheat (332 kg ha⁻¹) and *moong*-wheat (330 kg ha⁻¹) cropping systems. Sharma and Jain (2014) also reported highest available K status and actual K status in *guar*-wheat cropping system compared to maize-wheat and groundnut-wheat system.

Economics

The economic analysis of cropping system as a whole (Table 6) revealed that the highest cost of cultivation (Rs.70901 ha⁻¹) was recorded in cotton-wheat cropping sequence followed by cotton-*raya* (Rs.62819 ha⁻¹) and lowest was obtained under *guar*-barley (Rs.29690 ha⁻¹). The gross returns were also highest in cotton-wheat (Rs. 97192 ha⁻¹) followed by *guar*-wheat (Rs.90063 ha⁻¹) cropping pattern. Nevertheless, the net returns were lower in cotton-wheat than *guar*-wheat crop sequence. Jain *et al.* (2015) also reported the lowest input cost and higher (3.18:1) benefit: cost ratio in *guar*-wheat cropping system mainly due to lower fertilizer requirement and lower cost of cultivation. Sharma and Jain (2014), also reported highest benefit : cost ratio (3.66) in clusterbean-wheat cropping system compared to maize-wheat (3.43) and groundnut-wheat (3.48) cropping sequences. Cotton crop requires higher dose of nitrogen, more number of pesticides sprays and other inputs like cotton picking besides costly *Bt* seed etc., which leads to higher variable cost. The lowest net returns were found in *moong*-barley crop sequence. It may be due to lower potential yield of both *moong* and barley. Chauhan (2011) also found *guar*-wheat cropping sequence more profitable due to higher gross and net return and low input cost among *guar*-gram and *guar*-mustard crop sequences. When compared the increase or decrease over existing crop sequence (cotton-wheat), *guar* based crop sequences showed a positive increase than cotton and *moong* based crop sequences in terms of rupees ha⁻¹. Disparity in economics of various crop sequences has also been reported by Gawai and Pawar (2006).

Based on net returns and cotton equivalent yield it may be inferred that, *guar*-wheat crop sequence was more

remunerative as compared to other rotations if there is assured marketing and lucrative price (more than Rs 3000 q⁻¹). The crop sequence may be practiced where there is scarcity of water, as it needs lesser irrigations (1-2) than cotton (5) and can also be helpful in the formulation of contingent plan depending upon availability of water. Furthermore small or marginal area if replaced with existing cotton-wheat crop sequence may help to conserve water resources as well help in soil sustainability.

REFERENCES

- Chauhan, R. P. S. 2011. Studies on the crop diversification of existing cropping system in North-West Rajasthan. *J. Prog. Agric.* **2** (2): 39-40.
- Gawai, P. P. and V. S. Pawar, 2006. Integrated nutrient management in sorghum (*Sorghum bicolor*)-chickpea (*Cicer arietinum*) cropping sequence under irrigated conditions. *Indian J. Agron.* **51**: 17-20.
- Gomez, K.A. and A. A. Gomez, 1984. Statistical procedures for agricultural research. John Wiley & Sons, New Delhi.
- Jain, N. K., H. Singh, L. N. Dashora and S. L. Mundra, 2015. Maize (*Zea mays*)-wheat (*Triticum aestivum*) cropping system: Intensification through introduction of pulses. *Indian J. Agron.* **60** (3):347-351.
- Kadian, V. S., S. K. Thakral and S. Kumar, 2009. Evaluation of different crop sequences in south-west Haryana. *Haryana J. Agron.* **25** (1&2): 4-6.
- Sharma, S. K. and N. K. Jain, 2014. Nutrient management in wheat (*Triticum aestivum*)-based cropping systems in sub-humid southern zone of Rajasthan. *Indian J. Agron.* **59** (1): 26-33.
- Sidhu, R. S., Sukhpal Singh and A.S. Bhullar, 2011. Farmers' Suicides in Punjab: A census survey of the two most affected districts. *Eco. and pol. weekly.* **XLVI** (26 & 27):131-37.
- Singh, Sukhpal., H. S. Kingra, R. Singh, 2013. Input utilization and constraints of cotton production in Punjab. *J. Cotton Res. Dev.* **27** (1) : 144-148.
- Singh, N.B. and K. K. Verma, 1998. Production potential and economic analysis of rice (*Oryza sativa*) based cropping system. *Indian J. Agron.* **43**:199-203.
- Usadadiya, V. P. and R. H. Patel, 2013. Influence of preceding crops and nutrient management on productivity of wheat (*Triticum aestivum*)-based cropping system. *Indian J. Agron.* **58**(1):15-18.
- Varvel, G. E. and W.W. Wilhelm, 2003. Soybean nitrogen contribution to corn and sorghum in western corn belt rotations. *Agron. J.* **95**:1220-25.

Rec. on 10.05.2017 & Acc. on 12.06.2017