

# Journal of Soils and Crops

Volume 32

Number 1

June 2022

## C O N T E N T S

|                                                                                                                                                                                                                                                                                                                                                      |              |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|
| <b>Domestication and popularization of underutilised and wild fruits for fruit diversification : A review</b><br>Ritika Mehta <sup>1</sup> , Bornali Gogoi <sup>2</sup> , Himadri Shekhar Datta <sup>3</sup> and Purnima Pathak <sup>4</sup>                                                                                                         | <b>1-8</b>   |
| <b>In-silico identification of microRNA-like RNAs and their regulating target functions in <i>Fusarium graminearum</i>, a head blight pathogen of wheat</b><br>Lalit L. Kharbikar <sup>1</sup> , Arti S. Shanware <sup>2</sup> and Simon G. Edwards <sup>3</sup>                                                                                     | <b>9-18</b>  |
| <b>Isolation, identification and molecular characterisation of chromium tolerant bacteria from industrial effluent and their efficacy study</b><br>Vishnupriya Kondapalli and B. Sujatha                                                                                                                                                             | <b>19-24</b> |
| <b>Genetic recombination and gene amplification of <i>nif H</i> from free nitrogen fixer <i>Azotobacter chroococcum</i> to <i>Escherichia coli</i> by transformation</b><br>Vinay V. Chougule <sup>1</sup>                                                                                                                                           | <b>25-29</b> |
| <b>Characterization and classification of farm soils of regional agricultural research station (RARS), Nandyal, Kurnool District, Andhra Pradesh</b><br>S. Balaji Nayak <sup>1</sup> , S. Jaffar Basha <sup>2</sup> , M. Jyostna Kiranmai <sup>3</sup> and T. Bhagavatha Priya <sup>4</sup>                                                          | <b>30-36</b> |
| <b>Effect of row spacing, variety and foliar spray on quality, nutrient content and uptake of summer green gram (<i>Vigna radiata</i> L.) under South Gujarat condition</b><br>P. T. Patel <sup>1</sup> , A. D. Raj <sup>2</sup> and V. R. Jinjala <sup>3</sup>                                                                                      | <b>37-44</b> |
| <b>Micronutrient status and their relationship with leaf nutrient content of cauliflower (<i>Brassica oleracea</i> var. botrytis) growing areas of Kunah river flood plains in Hamirpur, Himachal Pradesh</b><br>Anil Kumar <sup>1</sup> , Mohammad Arshad <sup>2</sup> , Shiv Pratap Singh <sup>3</sup> and Swapana Sepehya <sup>4</sup>            | <b>45-58</b> |
| <b>Effect of sowing methods and row ratio on agronomic advantages, competition functions and economics of wheat (<i>Triticum aestivum</i> L.) + Chickpea (<i>Cicer arietinum</i> L.) intercropping</b><br>Ankushdeep and Rakesh Kumar                                                                                                                | <b>59-63</b> |
| <b>Effect of different application methods of nitrogen at seedling stage and different top dressing methods on yield of rice (<i>Oryza sativa</i> L.)</b><br>Manish Pradhan <sup>1</sup> , Jamkhogin Lhungdim <sup>2</sup> , Edwin Luikham <sup>3</sup> , A. Herojit Singh <sup>4</sup> , Pawan kumar <sup>5</sup> and N. Okendro Singh <sup>6</sup> | <b>64-70</b> |
| <b>Highlighting the potentialities of five underutilised spices used by the Karbi tribe of Assam, India</b><br>Moromi Engtipi <sup>1</sup> and N.L. Raju <sup>2</sup>                                                                                                                                                                                | <b>71-75</b> |
| <b>Study of soil stabilization using soda-lime and coir husk ash in Karewa soils</b><br>Minsa Ali <sup>1</sup> and Ashish Kumar <sup>2</sup>                                                                                                                                                                                                         | <b>76-79</b> |
| <b>Effect of nitrogen, phosphorus and potassium on growth, yield and economics of hybrid maize (<i>Zea mays</i> L.) under south Gujarat condition</b><br>K. V. Ram <sup>1</sup> , A. D. Raj <sup>2</sup> and P. M. Sankhla                                                                                                                           | <b>80-86</b> |
| <b>Intermolecular interaction of triple super phosphate fertilizer in aqueous soil salt solutions : an ultrasonic study</b><br>Paritosh L. Mishra <sup>1</sup> , Ajay B. Lad <sup>2</sup> and Urvashi P. Manik <sup>3</sup>                                                                                                                          | <b>87-91</b> |
| <b>Preferential flow : A hot spot for biological activity in soils - A case study</b><br>Pushpanjali <sup>1</sup> , A. Angayarkanni <sup>2</sup> , Ashish S Dhimate <sup>3</sup> , Josily Samuel <sup>4</sup> , A. G. K. Reddy <sup>5</sup> and Sannaila Arun Kumar <sup>6</sup>                                                                     | <b>92-96</b> |

|                                                                                                                                                                                                                                                                                                                                         |                |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <b>Effect of foliar application of potassium nitrate and salicylic acid on yield, yield attributes and economics of wheat (<i>Triticum aestivum</i> L.)</b><br>Rahul Monga <sup>1</sup> and Rakesh Kumar <sup>2</sup>                                                                                                                   | <b>97-101</b>  |
| <b>Influence of planting densities and nutrient levels on sweet corn genotypes in North Eastern Himalayan Region of India</b><br>Sanjenbam Dayananda Singh <sup>1</sup> , Jeti Konsam <sup>2</sup> , Nabakishor Nongmaithem <sup>3</sup> , Pushparani Senjam <sup>4</sup> and Diana Samurailatpam <sup>5</sup>                          | <b>101-105</b> |
| <b>Testing pathogenicity of white grub, <i>Holotrichia serrata</i> (Fabricius) (Coleoptera : Scarabaeidae) from the Darjeeling foothill tea plantation of Bengal</b><br>S. K. Subba <sup>1</sup> , R. Biswa <sup>2</sup> and M. Sarker <sup>3</sup>                                                                                     | <b>106-111</b> |
| <b>Influence of different slow release and controlled release fertilizers on yield and nutrient dynamics in soybean production</b><br>S. M. Todmal <sup>1</sup> , H. K. Kausadikar <sup>2</sup> , Syed Ismail <sup>3</sup> and S. L. Waikar <sup>4</sup>                                                                                | <b>112-122</b> |
| <b>Evaluation of rice genotypes for bio chemical and yield and yield contributing characters</b><br>Ajay F. Tangle <sup>1</sup> , G. R. Shamkuwar <sup>2</sup> , R. D. Deotale <sup>3</sup> and S.G. Shamkuwar <sup>4</sup>                                                                                                             | <b>123-126</b> |
| <b>Effect of foliar application of iron and zinc on morpho-physiological parameters and yield in safflower (<i>Carthamus tinctorius</i> L.)</b><br>Avinash P. Kesarkar <sup>1</sup> , P. V. Shende <sup>2</sup> , Vinita S. Karad <sup>3</sup> , P. R. Manapure <sup>4</sup> , R. D. Deotale <sup>5</sup> and S. D. Tupkar <sup>6</sup> | <b>127-132</b> |
| <b>Evaluation of chitosan concentrations for improving chemical, biochemical and yield contributing parameters and yield of chickpea</b><br>Ayushi N. Raut <sup>1</sup> , R.D. Deotale <sup>2</sup> , R.O. Gopale <sup>3</sup> , S. B. Baviskar <sup>4</sup> and V. S. Madke <sup>5</sup>                                               | <b>133-140</b> |
| <b>Impact of fertilization on uptake of nutrients, nutritional quality and yield of chickpea grown on farmers fields in Chandur Bazar Taluka of Amravati district</b><br>Shradha V. Bhule <sup>1</sup> , R. M. Ghodpage <sup>2</sup> , S. S. Balpande <sup>3</sup> and A.R. Mhaske <sup>4</sup>                                         | <b>141-145</b> |
| <b>Genetic analysis in F2 populations of linseed (<i>Linum usitatissimum</i> L.)</b><br>Yogesh P. Gite <sup>1</sup> , Prema R. Manapure <sup>2</sup> , Prashant V. Shende <sup>3</sup> , Shanti R. Patil <sup>4</sup> , Beena M. Nair <sup>5</sup> and Aambadas R. Aade <sup>6</sup>                                                    | <b>146-149</b> |
| <b>Effect of nutrient management practices on sulphur dynamics under chickpea grown in farmers field of Indapur taluka</b><br>Akshay D. Kate <sup>1</sup> , R.M. Ghodpage <sup>2</sup> , A.R. Mhaske <sup>3</sup> , Kajal D. Bhojar <sup>4</sup> and Bharat P. Rathod <sup>5</sup>                                                      | <b>150-154</b> |
| <b>Response of maize to silicon application</b><br>S.S.Mane <sup>1</sup> , K. T.Jadhav <sup>2</sup> and P. P. Markande <sup>3</sup>                                                                                                                                                                                                     | <b>155-158</b> |
| <b>Evaluation of white and pigmented rice cultivars for morpho-physiological characters and yield</b><br>Krunal N. Bele <sup>1</sup> , Gautam R. Shamkuwar <sup>2</sup> , Rajesh D. Deotale <sup>3</sup> , Aditi S. Deshmukh <sup>4</sup> and Sahil G. Shamkuwar <sup>5</sup>                                                           | <b>159-163</b> |
| <b>Analysis of soil to ensure NPK proportion at various sites of Panewadi village, District Nashik (M.S.) India</b><br>Balaji Pandurang More <sup>1</sup>                                                                                                                                                                               | <b>164-166</b> |
| <b>Cost of milk production of milch buffaloes maintained in nilanga tahsil of Latur district</b><br>Priyanka Katke <sup>1</sup> , V. G. Atkare <sup>2</sup> and N. D. Kapgate <sup>3</sup>                                                                                                                                              | <b>167-170</b> |
| <b>Effectivity of different sowing dates on chemical, biochemical, yield and yield traits in mustard genotypes</b><br>Rahul Gopale <sup>1</sup> , R. D. Deotale <sup>2</sup> , Ayushi Raut <sup>3</sup> , S. R. Kamdi <sup>4</sup> , S. B. Baviskar <sup>5</sup> and Ambadas Aade <sup>6</sup>                                          | <b>171-179</b> |

|                                                                                                                                                                 |                |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <b>Estimation of heterosis for yield and yield contributing traits in linseed (<i>Linum usitatissimum</i> L.)</b>                                               | <b>180-183</b> |
| Suvarnlata D. Shivankar <sup>1</sup> , Beena M. Nair <sup>2</sup> and Dhananjay S. Thikare <sup>3</sup>                                                         |                |
| <b>Effect of foliar application of micronutrients on chemical, biochemical parameters and yield attributes in chickpea (<i>Cicer arietinum</i> L.)</b>          | <b>184-192</b> |
| Vinita Sudam Karad <sup>1</sup> , P. V. Sihende <sup>2</sup> , A. P. Kesarkar <sup>3</sup> , V. S. Madake <sup>4</sup> and S. D. Tupkar <sup>5</sup>            |                |
| <b>Impact of stress irrigation management through partial root-zone drying on yield and soil parameters of tomato (<i>Lycopersicon esculentum</i> Mill.)</b>    | <b>193-199</b> |
| Komal Chavan <sup>1</sup> , P. S. Bodake <sup>2</sup> , V.G. Raut <sup>3</sup> and Darshana Kadam <sup>4</sup>                                                  |                |
| <b>Economics of different integrated farming system models, under irrigated conditions of Western Maharashtra</b>                                               | <b>200-207</b> |
| P. P. Kharche <sup>1</sup> , U. S. Surve <sup>2</sup> , A. D. Tumbare <sup>3</sup> and V. G. Raut <sup>4</sup>                                                  |                |
| <b>Studies on involvement of rural woman in preparation of milk and milk products in Tirora Tahsil of Gondia District (M.S.)</b>                                | <b>208-211</b> |
| Rubi Katre <sup>1</sup> , V. G. Atkare <sup>2</sup> , N. S. Chore <sup>3</sup> and Pranav Adpawar <sup>4</sup>                                                  |                |
| <b>Seed viability and vigour assessment of different seed lots of tetraploid and diploid cotton</b>                                                             | <b>212-216</b> |
| Radhesham V. Hatzade <sup>1</sup> , Sunil S. Mahajan <sup>2</sup> , Eugenia P. Lal <sup>3</sup> , Suchit A. John <sup>4</sup> and Pradip K. Shukla <sup>5</sup> |                |