

## EFFICACY OF PUTRESCINE AND IBA ON BIOCHEMICAL AND YIELD CONTRIBUTING PARAMETERS OF BLACK GRAM

R. D. Deotale<sup>1</sup>, N. D. Jadhav<sup>2</sup>, Shanti Patil<sup>3</sup>, Sapana Baviskar<sup>4</sup>,  
Vandana S. Madke<sup>5</sup> and Vandana Kalamkar<sup>6</sup>

### ABSTRACT

In order to examine effects of different concentrations of growth regulators (putrescine and IBA) on biochemical and yield contributing parameters on black gram a field experiment was carried out in the Botany farm, College of Agriculture, Nagpur during 2017-18. Foliar application of putrescine and IBA 25, 50, 75 and 100 ppm each was given at vegetative stage (25 and 40 DAS) and the data were taken at 25, 40 and 55 DAS coinciding with vegetative stage, flowering stage and before harvest stage. Application of putrescine and IBA enhanced biochemical parameters (*viz.*) chlorophyll content, NPK content in leaves and protein content in grains, yield contributing characters *viz.*, number of seeds pod<sup>-1</sup>, weight of 100 seeds, seed yield plant<sup>-1</sup>, plot<sup>-1</sup> and ha<sup>-1</sup>. Analysis of data revealed that 100 ppm IBA considered as a most effective concentration in enhancing all biochemical and yield contributing parameters. But considering the B : C ratio two foliar sprays of 25 ppm IBA at 25 and 40 DAS was found more effective and economical treatment with increased yield of 30.47% having B:C ratio of 2.97 as compared to 2.54 in control.

(Key words : black gram, putrescine, IBA, foliar application, biochemical parameters and yield)

### INTRODUCTION

Black gram is chief constituent of dal (husked or unhusked), papad, idali, dahi wada, and imarati. The grain of black gram is superior in nutritive value, black gram contain about 24% proteins, 61% carbohydrate, 1.8% fat, 345 Kcal energy, 3.5% minerals, 385 mg 100 g<sup>-1</sup> phosphoric acid and 16.2% total dietary fibre. Black gram rich in minerals like Zinc, Calcium and also rich in Vitamins like thiamine, riboflavin and niacin.

In India, black gram traditionally grown in *kharif* season, but in south it also grown as *rabi* crop. It is widely grown in India and Central Asia. In India generally grown in area of which received annual rainfall of 800 mm. It is a hardy and drought resistant plant. Major black gram producing state in India is Maharashtra. Maharashtra is a leading producer of black gram followed by Andhra Pradesh, Uttar Pradesh, Madhya Pradesh, Tamilnadu, Karnataka, Rajasthan and Orissa.

The yield of black gram may be enhanced through physiological manipulation such as foliar application of

putrescine and IBA. Plant growth regulators are shown to change leaf resistance by altering stomatal aperture, the rate of photosynthesis could be manipulated through this technology. Putrescine, IBA, IAA, GA, kinetin, phenolics and aliphatic alcohols are reported to increase and stimulate the rate of photosynthesis. The diamine putrescine occurred widely in the higher plants. It was suggested to be involved in a variety of growth and developmental processes such as cell division (Bueno and Matilla, 1992), fruit set and growth (Biasi *et al.*, 1991) and senescence (Kao, 1994).

Considering the above facts present investigation was undertaken to study the effect of putrescine and IBA on biochemical and yield and yield contributing parameters of black gram.

### MATERIALS AND METHODS

A field experiment was conducted at Botany farm, College of Agriculture, Nagpur to know the response of black to growth regulators on its biochemical and yield contributing parameters. The experiment was laid out in

- 
1. Professor, Botany Section, College of Agriculture, Nagpur
  2. P.G. Student, Botany Section, College of Agriculture, Nagpur
  - 3, 4 and 5. Asstt. Professors, Botany Section, College of Agriculture, Nagpur
  6. Senior Research Assistant, Botany Section, College of Agriculture, Nagpur

complete randomized block design with three replications. The experiment consisted of nine treatments viz., IBA and putrescine applied at 25, 50, 75 and 100 ppm each with control (water spray). The seeds were sown with a spacing of 30 cm x 10 cm. Growth promoters were sprayed twice at 25 and 40 DAS. First observation (25 DAS) was made before the spray and other observations were made at 40 and 55 DAS. Five randomly selected plants were tagged and observations were taken on biochemical parameters (leaf chlorophyll content, N, P, K content in leaves and protein content) and yield attributing characters (Number of seeds pod<sup>-1</sup>, weight of 100 seeds and yield plot<sup>-1</sup>). Determination of nitrogen and protein was carried out by micro-kjeldhal method as given by Somichi *et al.* (1972). Phosphorus and potassium estimated by vanadomolybdate yellow colour method, (using calorimeter and flame photometer respectively) given by Jackson (1967). The observed data were analyzed statistically using analysis of variance at 5% level of significance (Panse and Sukhamate, 1954).

## RESULTS AND DISCUSSION

### Chlorophyll content

The chlorophyll content differed significantly between control and growth regulators applied plants (Table 1). The content varied with the lowest value of 1.28 mg g<sup>-1</sup> in control (T<sub>1</sub>) to highest value of 1.94 mg g<sup>-1</sup> in 100 ppm IBA treatment (T<sub>9</sub>) at 40 DAS and lowest value of 1.19 in control (T<sub>1</sub>) to highest value of 1.49 in 100 ppm IBA treatment (T<sub>9</sub>) at 55 DAS among the treatments studied. Foliar application of all the growth regulator treatments increased the chlorophyll content over control except application of 75 ppm putrescine (T<sub>4</sub>), 50 ppm IBA (T<sub>7</sub>), 50 ppm putrescine (T<sub>3</sub>), 25 ppm IBA (T<sub>6</sub>) and 25 ppm putrescine (T<sub>2</sub>). These treatments were found at par with control. Among the treatments, the highest chlorophyll content was recorded in treatment T<sub>9</sub> (100 ppm IBA) followed by the treatments T<sub>5</sub> (100 ppm putrescine) and T<sub>8</sub> (75 ppm IBA) when compared with treatment T<sub>1</sub> (control) at both the stages of observations. Putrescine or IBA treatments retard senescence and chlorophyll loss by altering the stability and permeability of membranes and protecting chloroplast from senescing (Gonzalez-Aguilar *et al.*, 1997). A possible explanation for the promoting effect of putrescine on photosynthetic pigment of black gram plant in the present work is that PAs might retard the chlorophyll destruction and / or increase their biosynthesis or stabilize the thylakoid membrane. Similar effects of PAs on photosynthetic pigments had been observed by Pinkey *et al.* (2016). They conducted a field experiment to access the effect of putrescine in mitigating the water stress in wheat and reported that foliar sprays of 50 and 100 ppm putrescine significantly increased the chlorophyll content.

### Nitrogen content

Leaf nitrogen content differed significantly among the treatments and varied from a minimum of 3.96% at 40 DAS and 3.56% at 55 DAS in control treatment to a maximum

of 5.81% at 40 DAS and 4.77% at 55 DAS in treatment T<sub>9</sub> (100 ppm IBA). Next to this treatment the treatments were T<sub>5</sub> (100 ppm putrescine), T<sub>8</sub> (75 ppm IBA) and T<sub>4</sub> (75 ppm putrescine) when compared with treatment T<sub>1</sub> (control). Whereas, treatments T<sub>3</sub> (50 ppm putrescine), T<sub>6</sub> (25 ppm IBA) and T<sub>2</sub> (25 ppm putrescine) were found at par with treatment T<sub>1</sub> (control).

From this data it is observed that leaf nitrogen content was increased up to 40 DAS and reduced thereafter, at 55 DAS. The decrease in nitrogen content might be due to fact that younger leaves and developing organs, such as grains act as strong sink demand and may draw heavily nitrogen from older leaves (Gardner *et al.*, 1988). Results recorded by Poonkodi (2003) also stated that decrease in nitrogen content at later stage might be due to translocation and utilization of nutrient for flower and pod formation.

Ahmed *et al.* (2013) tried putrescine and humic acid on cotton. These growth regulators were sprayed eight times started at 45 DAP and repeated every after 15 days on cotton and reported that foliar application of 2 ppm putrescine and 1% humic acid significantly enhanced inorganic nitrogen.

### Phosphorus content

Leaf phosphorus content at 40 DAS differed among the treatments and varied from a minimum of 0.833% in control to a maximum of 0.938% in treatment T<sub>9</sub> (100 ppm IBA). Treatment T<sub>5</sub> (100 ppm putrescine) also showed their significance and enhanced phosphorus content in leaves when compared with treatment control and other treatments under study. While, foliar application of 75 ppm IBA (T<sub>8</sub>), 75 ppm putrescine (T<sub>4</sub>), 50 ppm IBA (T<sub>7</sub>) and 50 ppm putrescine (T<sub>3</sub>) also increased phosphorus content significantly over control. Treatments T<sub>6</sub> (25 ppm IBA) and T<sub>2</sub> (25 ppm putrescine) were found at par with control (T<sub>1</sub>).

Leaf phosphorus content at 55 DAS differed among the treatments and varied from a minimum of 0.812% to a maximum of 0.911% in treatment T<sub>9</sub> (100 ppm IBA) followed by the treatments T<sub>5</sub> (100 ppm putrescine) T<sub>8</sub> (75 ppm IBA) and T<sub>4</sub> (75 ppm putrescine) when compared with treatment T<sub>1</sub> (control) and remaining treatments under study. But treatments T<sub>7</sub> (50 ppm IBA), T<sub>3</sub> (50 ppm putrescine), T<sub>6</sub> (25 ppm IBA) and T<sub>2</sub> (25 ppm putrescine) were found at par with control in leaf phosphorus content (Table 1).

Deotale *et al.* (2017) observed that foliar sprays of putrescine and IBA @ 100 ppm significantly enhanced phosphorus content in soybean leaves.

### Potassium content

Potassium is an essential macronutrient for plants involved in many physiological processes. It is important for crop yield as well as for the quality of edible parts of crops. Although potassium is not assimilated into organic matter, potassium deficiency has a strong impact on plant metabolism. Plant responses to low potassium involve changes in the concentrations of many metabolites as well as alteration in the transcriptional levels of many genes and in the activity of many enzymes.

Data pertaining to potassium content in leaves were estimated at various stages of observations viz., 25, 40 and 55 DAS. Significant results were recorded at all the stages of observations viz., 40 and 55 DAS except 25 DAS.

At 40 DAS significantly highest leaf potassium content was observed in treatment T<sub>9</sub> (100 ppm IBA) followed by treatments T<sub>5</sub> (100 ppm putrescine), T<sub>8</sub> (75 ppm IBA) and T<sub>4</sub> (75 ppm putrescine). Similarly treatments T<sub>7</sub> (50 ppm IBA) and T<sub>3</sub> (50 ppm putrescine), were also found significantly superior over treatment T<sub>1</sub> (control) in potassium content. Treatments T<sub>6</sub> (25 ppm IBA) and T<sub>2</sub> (25 ppm putrescine) were found at par with treatment T<sub>1</sub> (control). The range of potassium content at 40 DAS in black gram was 0.98-1.91%.

At 55 DAS significantly highest leaf potassium content was observed in treatment T<sub>9</sub> (100 ppm IBA) followed by treatments T<sub>5</sub> (100 ppm putrescine) and T<sub>8</sub> (75 ppm IBA). Similarly treatments T<sub>4</sub> (75 ppm putrescine), T<sub>7</sub> (50 ppm IBA) and T<sub>3</sub> (50 ppm putrescine) were also found significantly superior over treatment T<sub>1</sub> (control). Treatments T<sub>6</sub> (25 ppm IBA) and T<sub>2</sub> (25 ppm putrescine) were found at par with treatment T<sub>1</sub> (control). The range of potassium content at 55 DAS in black gram was 0.82-1.62%.

It is clear from the data, that leaf potassium content was decreased gradually in later stage of crop growth i.e. 55 DAS. It might be because of diversion of potassium towards developing parts i.e. pod formation in black gram.

Deotale *et al.* (2016) conducted a field experiment to investigate the foliar sprays of putrescine and IBA on soybean and noted that two foliar sprays of 100 ppm putrescine and 100 ppm IBA at two stages i.e. before flowering and 10 days after flowering were found to be most effective in enhancing potassium content in leaves.

#### Protein content in seed

Significant variation in protein content was observed among the treatments studied (Table 2). Foliar application of growth regulator 100 ppm IBA registered the highest protein content (26.50%) followed by treatments T<sub>5</sub> (100 ppm putrescine), T<sub>8</sub> (75 ppm IBA), T<sub>4</sub> (75 ppm putrescine) and T<sub>7</sub> (50 ppm IBA) when compared with treatment T<sub>1</sub> (control) and rest of the treatments under study. Similarly, treatments T<sub>3</sub> (50 ppm putrescine) and T<sub>6</sub> (50 ppm IBA) were also increased seed protein significantly over treatment T<sub>1</sub> (control). Treatment T<sub>2</sub> (25 ppm putrescine) was found at par with treatment T<sub>1</sub> (control) in protein content.

It is observed from the results that putrescine and IBA are more effective in seed protein content. This might be due to enhancement of enzymatic activity and translocation of metabolites to the black gram seeds.

Mathur and Vyas (2007) conducted an experiment to study the effect of salicylic acid, sisterol as well as putrescine on pearl millet (*Pennisetum thyphoides*). Results showed that application of salicylic acid at 3 mM and sisterol or putrescine at 0.15 Mm significantly increased protein content.

#### Yield contributing parameters

Results presented in table 2 showed that both growth regulators (putrescine and IBA) have a promoting effect on yield contributing characters viz., number of seeds pod<sup>-1</sup>, 100 seed weight, yield plant<sup>-1</sup> and yield plot<sup>-1</sup> of black gram. Significant variation in seed weight was noticed among the treatments. The highest 100 seed weight (4.93 g) and the lowest 100 seed weight (4.10 g) were recorded in 100 ppm IBA (T<sub>9</sub>) and control (T<sub>1</sub>) respectively. However, 100 seed weight was significantly maximum in treatments T<sub>5</sub> (100 ppm putrescine), T<sub>8</sub> (75 ppm IBA), T<sub>4</sub> (75 ppm putrescine) and T<sub>7</sub> (50 ppm IBA) when compared with treatment T<sub>1</sub> (control). But treatments T<sub>3</sub> (50 ppm putrescine), T<sub>6</sub> (25 ppm IBA) and T<sub>2</sub> (25 ppm putrescine) were found at par with treatment T<sub>1</sub> (control).

Number of seeds pod<sup>-1</sup> varied among the treatments and ranged a minimum of 4.51 to a maximum of 6.18. Among all the treatments the highest seeds pod<sup>-1</sup> was obtained in treatment T<sub>9</sub> (100 ppm IBA) followed by treatments T<sub>5</sub> (100 ppm putrescine), T<sub>8</sub> (75 ppm IBA) and T<sub>4</sub> (75 ppm putrescine) when compared with treatment T<sub>1</sub> (control) and rest of the treatments under observations. Foliar application of 50 ppm putrescine, 25 ppm IBA and 25 ppm putrescine exhibited significantly minimum number of seeds pod<sup>-1</sup> and remained at par with control treatment.

Seed yield plant<sup>-1</sup> varied among the treatments. Foliar application of growth regulators significantly increased seed yield plant<sup>-1</sup> over control. The highest seed yield (4.05 g plant<sup>-1</sup>) was recorded in treatment T<sub>9</sub> (100 ppm IBA) as compared to (2.36 g plant<sup>-1</sup>) treatment T<sub>1</sub> (control). Seed yield plot<sup>-1</sup> also varied among the treatments. Foliar application of growth regulators significantly increased seed yield plot<sup>-1</sup> over control. The highest seed yield (0.57 kg plot<sup>-1</sup>) was recorded in treatment T<sub>9</sub> (100 ppm IBA) as compared to (0.33 kg plot<sup>-1</sup>) treatment T<sub>1</sub> (control). Deotale and Pandey (2017) also reported that foliar application of 100 ppm IBA significantly increased grain yield of maize. Shinde *et al.* (2018) reported that foliar application 100 ppm IBA followed by 100 ppm putrescine significantly increased pod yield of groundnut.

The highest seed yield of 13.49 q ha<sup>-1</sup> was obtained by the treatment T<sub>9</sub> (100 ppm IBA) which registered 52.25% higher seed yield over control (8.86 q ha<sup>-1</sup>). Treatments T<sub>5</sub> (100 ppm putrescine), T<sub>8</sub> (75 ppm IBA), T<sub>4</sub> (75 ppm putrescine), T<sub>7</sub> (50 ppm IBA), T<sub>3</sub> (50 ppm putrescine), T<sub>6</sub> (25 ppm IBA) and T<sub>2</sub> (25 ppm putrescine) in a descending manner also enhanced seed yield ha<sup>-1</sup> significantly over control.

The increase in yield and yield attributes may be due to altering the hormonal balance and improved water relation in plants. PA namely putrescine is involve in stabilization of D<sub>1</sub> and D<sub>2</sub> polypeptides of photosystem second which is the source of electron for NADP<sup>+</sup> reduction at photosystem one (Taiz and Zeiger, 1991). It is also prevented the lipid peroxidation, proteolytic attack and inhibitors of ethylene synthesis through inhibition of ACC synthase and conversion of ACC to ethylene, which is a

Table 1. Effect of putrescine and IBA on leaf chlorophyll, nitrogen and phosphorus contents

Treatment	Chlorophyll content in leaves (mg g <sup>-1</sup> )			Nitrogen content in leaves (%)			Phosphorous content in leaves (%)		
	25 DAS	40 DAS	55 DAS	25 DAS	40 DAS	55 DAS	25 DAS	40 DAS	55 DAS
T <sub>1</sub> (control)	1.22	1.28	1.19	3.48	3.96	3.56	0.632	0.833	0.812
T <sub>2</sub> (25 ppm putrescine)	1.28	1.35	1.21	3.70	4.12	3.74	0.653	0.847	0.826
T <sub>3</sub> (50 ppm putrescine)	1.24	1.46	1.33	3.49	4.59	4.15	0.678	0.869	0.842
T <sub>4</sub> (75 ppm putrescine)	1.32	1.58	1.38	3.82	5.19	4.63	0.629	0.892	0.876
T <sub>5</sub> (100 ppm putrescine)	1.29	1.78	1.47	3.57	5.52	4.87	0.664	0.921	0.897
T <sub>6</sub> (25 ppm IBA)	1.23	1.42	1.24	3.73	4.22	3.83	0.667	0.856	0.834
T <sub>7</sub> (50 ppm IBA)	1.34	1.52	1.34	3.66	4.72	4.27	0.618	0.878	0.857
T <sub>8</sub> (75 ppm IBA)	1.26	1.65	1.46	3.83	5.34	4.77	0.646	0.903	0.893
T <sub>9</sub> (100 ppm IBA)	1.33	1.94	1.49	3.67	5.81	5.17	0.658	0.938	0.911
SE (m) ±	0.08	0.11	0.07	0.21	0.23	0.27	0.04	0.01	0.02
CD at 5%	-	0.32	0.20	-	0.68	0.80	-	0.03	0.05

**Table 2. Effect putrescine and IBA on leaf potassium content, seed protein, seeds pod<sup>-1</sup>, 100 seed weight, seed yield plant<sup>-1</sup>, plot<sup>-1</sup>, ha<sup>-1</sup>, per cent increase in yield and B:C ratio**

Treatments	Leaf potassium content (%)			Protein Content (%)	Seeds pod <sup>-1</sup>	100 seed weight (g)	Seed yield plant <sup>-1</sup> (g)	Seed yield plot <sup>-1</sup> (kg)	Seed yield ha <sup>-1</sup> (q)	Per cent increase over control	B:C ratio
	25 DAS	40 DAS	55 DAS								
T <sub>1</sub> (control)	0.81	0.98	0.82	24.20	4.51	4.10	2.36	0.33	8.86	-	2.54
T <sub>2</sub> (25 ppm putrescine)	0.87	1.05	0.89	24.86	4.70	4.23	3.27	0.46	10.89	22.91	2.37
T <sub>3</sub> (50 ppm putrescine)	0.92	1.46	1.07	25.54	5.22	4.50	3.62	0.51	12.06	36.11	2.11
T <sub>4</sub> (75 ppm putrescine)	0.84	1.69	1.32	25.98	5.61	4.66	3.79	0.53	12.63	42.50	1.85
T <sub>5</sub> (100 ppm putrescine)	0.89	1.88	1.49	26.25	6.05	4.80	4.03	0.56	13.43	51.58	1.69
T <sub>6</sub> (25 ppm IBA)	0.86	1.24	0.98	25.32	4.98	4.30	3.47	0.49	11.56	30.47	2.97
T <sub>7</sub> (50 ppm IBA)	0.91	1.54	1.22	25.68	5.31	4.65	3.69	0.52	12.29	38.71	2.86
T <sub>8</sub> (75 ppm IBA)	0.85	1.74	1.42	26.10	5.74	4.70	3.85	0.54	12.83	44.80	2.73
T <sub>9</sub> (100 ppm IBA)	0.94	1.91	1.62	26.50	6.18	4.93	4.05	0.57	13.49	52.25	2.64
SE (m) ±	0.06	0.09	0.07	0.31	0.24	0.14	0.21	0.03	0.56	-	-
CD at 5%	-	0.26	0.20	0.92	0.71	0.41	0.62	0.08	1.67	-	-

common phenomenon occurred during senescence. The present results also in line with the report of Kabir *et al.*, 1992. They stated that PAs improved the yield and yield attributes in tomato. The increments in yield components due to putrescine treatments may be attributed to the increasing growth rate, in this respect, Davies (1995) reported that polyamines play a critical role in different biological processes, including cell division, growth, somatic embryogenesis, floral initiation, development of flowers and fruits. It is worthy to mention that there is a close relationship between the effect of PAs and the stimulated growth, endogenous phytohormones, the photosynthetic output (soluble sugars, polysaccharides and total carbohydrates) and the nitrogen constituents. These results might increase the efficiency of solar energy conversion which maximize the growth ability of black gram plant and consequently increased its productivity and yield components.

Amin *et al.* (2013) explained the effect of two plant growth regulators putrescine and Indole-3-butyric acid (IBA) @ 25, 50 and 100 mg l<sup>-1</sup> applied either alone or in combinations. Spraying of putrescine and IBA @100 mg l<sup>-1</sup> significantly increased number of pods, seed yield, straw and biological yield feed<sup>-1</sup> of chickpea (*Cicer arietinum*).

## REFERENCES

- Ahmed, A.H. Hanafy, E. Darwish, S.A. F. Hamoda and M.G. Alobaidy, 2013. Effect of putrescine and humic acid on growth, yield and chemical composition of cotton plants grown under saline soil conditions. *Am-Euras. J. Agric. & Environ. Sci.* **13** (4): 479-497.
- Amin, A. A., F.A. Gharib, H.F. Abouzienna and Mona G. Dawood, 2013. Role of indole-3-butyric acid or/and putrescine in improving productivity of Chickpea (*Cicer arietinum* L.) Plants. *Pakistan J. Biol. Sci.* **16**: 1894-1903.
- Biasi, R., G. Costa and N. Bagni, 1991. Polyamine metabolism as related to fruit set and growth. *Pl. Physiol. Biochem.* **29**:497-506.
- Bueno, M. and A. Matilla, 1992. Effect of spermine and abscisic acid on mitotic divisions in isolated embryonic axes of chick pea seeds. *Cytobiology*, **71**:151-155.
- Davies, P.J. 1995. *Plant hormones: Physiology and Biochemistry and Biology* Kluwer Academic Publishers, London. p.p. 159.
- Deotale, R.D., S.R. Patil, Y. A. Wagh, V. B. Kalamkar and V.S. Jayade, 2017. Influence of putrescine and indole-3-butyric acid on chemical and biochemical parameters and yield of soybean. *J. Soils and crops*, **27**(2) : 135-141.
- Deotale, R.D. and B.B. Pandey, 2017. An evaluation of potentiality of exogenously applied putrescine and IBA on biochemical and yield contributing parameters of maize, *J. Soil and Crop*, **27**(2) 47-53
- Gardner, F.P., R.B. Pearce, R.L. Mitchell, 1988. In: *Physiology of Crop Giants*. Iowa State University Press, Amen, Iowa, pp. 187-208.
- Gonzalez-Aguilar, G.A., L. Zacarias, M. Mulas and M.T. Lafuente, 1997. Temperature and duration of water dips influence chilling injury, decay and polyamine content in Fortune mandarins. *Postharvest Biol. Technol.* **12**: 61-69.
- Jackson, M. L. 1967. *Soil Chemical analysis*, Printice Hall of India Pvt. Ltd., New Delhi, pp. 25-28.
- Kabir, J., T. K. Maity and Partha Sarthy. 1992. Polyamine, ethylene and other physico-chemical parameters in tomato (*Lycopersicon esculentum*) fruits. *Sci. Cult.*, **58**: 159.
- Kao, C. H. 1994. Endogenous polyamine levels and dark- induced senescence of detached corn leaves. *Bot. Bull. Acad. Sin.* **35**:15-18.
- Mathur, Nishi and Anil Vyas, 2007. Physiological effect of some bioregulators on vegetative growth, yield and chemical constituents of pearl millet (*Pennisetum typhoides* (Burm) Stapf. and Hubb). *Int. J. Agri. Res.* **2** (3): 238-245.
- Panse, V.G. and P.V. Sukhante, 1954. *Statistical method for agriculture works*, ICAR New Delhi, p.p. 107-109.
- Pinkey Meena, N.S. Solanki and L. N. Dashora, 2016. Effect of putrescine on growth and productivity of wheat under water stress conditions. *Ann. Agric. Res. New Series*, **37** (1): 56-60.
- Poonkodi, P. 2003. Phosphorus use efficiency in black gram with pressmad. *Adv. Pl. Sci.* **17** (1): 239 -241.
- Shinde, R.D., R.D. Deotale and G.N. Jadhav, 2018. Role of putrescine and indole-3-butyric acid in improving morpho-physiological parameters and productivity of groundnut. *J. Soils and Crops*, **28** (1) : 81-88
- Somichi, Y., S. Y. Doughlus and A. P. James, 1972. *Laboratory manual. Physiological studies in rice analysis for total nitrogen (organic N) in plant tissue*. The inter. Res. Instti. Los Banos, Languna, Phillipine : II.
- Wagh, Y. A. 2015. Influence of putrescine and indole-3-butyric acid on growth and productivity of soybean. M.Sc. (Agri.) thesis (Unpublished) submitted to Dr. P.D.K.V., Akola.
- Taiz, L. and E. Zeiger. 1991. *Plant Physiology*. Benjamin/Cummings Publ. Company Inc. California.

**Rec. on 03.07.2019 & Acc. on 10.07.2019**