

EVALUATION OF MORPHO -PHYSIOLOGICAL TRAITS AND YIELD OF GROUNDNUT BY FOLIAR APPLICATION OF CHITOSAN

Kantilal B. Chande¹, Rajesh D. Deotale², Puja R. Mate³, Ajay A. Purane⁴ and Sapana B. Baviskar⁵

ABSTRACT

An investigation was carried out to study the effect of different concentrations of chitosan (control, 25, 50, 75, 100 and 125 ppm) in groundnut. The experiment was conducted at farm of Botany section, College of Agriculture, Nagpur during the *kharif* season of 2019-2020. Observations of plant height, number of branches, leaf area, dry matter were recorded at 25, 45, 65 and 85 DAS. RGR and NAR were calculated at 45-25, 65-45, 85-65 DAS. Yield plot⁻¹ was also recorded. Foliar sprays of chitosan at 25 and 45 DAS significantly enhanced above mentioned parameters with the application of 100 ppm chitosan when compared with other treatments and control.

(Key words: Groundnut, chitosan, foliar application, morpho-physiological parameters, yield)

INTRODUCTION

Groundnut, any of several plants that bear edible fruit or other nut like parts is the member of the family fabaceae or Leguminosae. It is a day neutral, self-pollinated crop plant having chromosome number $2n=40$. Groundnut belongs to family leguminosae, genus *Arachis* and has chromosome number $2n=40$. It is also a day neutral plant, flowers are self-pollinated.

Groundnut is cultivated as *kharif*, *rabi* and summer crop in India. Groundnut is mostly cultivated in South. During *kharif*-2018 all India groundnut acreage was 38, 90, 000 hectares. Five states, Gujarat 14,67,600 ha; 37.7%, Andhra Pradesh 6,60,000 ha; 17%, Rajasthan 5,49,052 ha; 14.1%, Karnataka 3,82,940 ha; 9.8%, Maharashtra 1,95,594 ha; 5.0% jointly accounted for 83.7% of the national acreage. At the national level, there was a decrease in acreage by 6.3% with respect to *kharif* 2017. The maximum decrease was observed for Gujarat 10.0% while, it was negligible for Andhra Pradesh 1.0%. The observed increase in acreage in Karnataka was nominal 1.3% (Anonymous, 2018).

Studies have found that chitosan oligosaccharides have the function of regulating plant development, and are a natural plant growth regulator. They can root and promote stems, shorten stems, and thrive, which is conducive to nutrients to maximize the supply of fruits. Chitosan has wide adaptability to plants, good affinity, no toxicity, no side effects, no harm to humans and animals, biode-

gradability, no soiling in the soil, slow degradation to small molecular carbohydrates, no pollution to the environment, an excellent pure natural ecological preparation. The positive effect of chitosan on plant growth may be credited to an increase in the key enzyme activities of nitrogen metabolism and increased photosynthesis which enhanced plant growth (Gornik *et al.*, 2008; Mondal *et al.*, 2012). Chitosan applications significantly increased total carbohydrates accumulation in the fruits (Mawgoud *et al.*, 2010). Hence, considering above facts an experiment was conducted to study the effect of different concentrations of chitosan on groundnut.

MATERIALS AND METHODS

An experiment was carried out at Shankar nagar farm of Agril. Botany section, College of Agriculture, Nagpur during 2019-20 in RBD with four replications and six treatments (control, 25, 50, 75, 100 and 125 ppm). Experimental gross plot was 3.15 m X 2.20 m and net plot was 2.10 m X 2.20 m. Observations on plant height, number of branches, plant⁻¹, leaf area, dry matter were recorded periodically at 25, 45, 65 and 85 DAS. RGR was calculated as per formula given by Blackman (1919) and NAR was calculated by using formula suggested by Williams (1946). The RGR and NAR were calculated at 45-25, 65-45 and 85-65 DAS. Yield plot⁻¹ was also recorded.

1, 3 & 4. P.G. Students, Agril. Botany Section, College of Agriculture, Nagpur

2. Professor, Agril. Botany Section, College of Agriculture, Nagpur

5. Asstt. Professor, Agril. Botany Section, College of Agriculture, Nagpur

RESULTS AND DISCUSSION

Plant height

Plant height is an important character of the vegetative phase and indirectly influences the yield components. It is the shortest vertical distance between the upper boundary of the main photosynthetic tissue on a plant and the stem or shoot base from the ground level.

At 45 DAS plant height was recorded significantly maximum in treatment T₅ (100 ppm chitosan) when compared with treatment T₁ (control) and remaining treatments under study. The treatments T₄ (75 ppm chitosan), T₃ (50 ppm chitosan), T₆ (125 ppm chitosan) and T₂ (25 ppm chitosan) were found at par with treatment T₁ (control) in plant height.

At 65 DAS highest plant height was recorded in treatment T₅ (100 ppm chitosan) when compared with treatment T₁ (control) and remaining treatments under study. Treatments T₄ (75 ppm chitosan), T₃ (50 ppm chitosan) and T₂ (25 ppm chitosan) were at par with each other but found significantly superior when compared with treatment T₁ (control) and remaining treatments. Treatments T₆ (125 ppm chitosan) and T₁ (control) were found at par with each other in plant height.)

At 85 DAS highest plant height was recorded in treatment T₅ (100 ppm chitosan) when compared with treatment T₁ (control) and remaining treatments under study. Furthermore treatments T₃ (50 ppm chitosan) and T₂ (25 ppm chitosan) significantly increased plant height over control and T₆ (125 ppm chitosan). Whereas, T₆ (125 ppm chitosan) also found significantly superior over control in plant height.

The per cent increase in plant height with respect to foliar application of 100 ppm chitosan were 17.00 at 45 DAS, 19.00 at 65 DAS and 42.24 at 85 DAS when compared with control (T₁).

Chitosan improved plant height might be due to increased number of internodes or length of internodes due to increased cell number (Hong Yan and Shu Yu, 2001).

Mondal *et al.* (2013) applied various concentrations of chitosan (0, 50, 25, 75 and 100 ppm) sprayed at 3 different stages viz., 35, 50, and 65 DAS. They suggested that foliar application of chitosan at 100 ppm expressively increased plant height in okra.

Number of the branches plant⁻¹

Branches are the site of the leaves, flower and peg formation. Hence, they are closely associated with the photosynthetic activity and yield of plant. So, number of branches is desirable attribute for higher biomass production and yield.

At 45 DAS impressive increase in number of branches was observed in treatment T₅ (100 ppm chitosan) followed by treatment T₄ (75 ppm chitosan) when compared with treatment T₁ (control) and rest of the treatments under study. Treatments T₃ (50 ppm) and T₂ (25 ppm) also found superior among rest of the treatments and treatment T₁

(control). Whereas, treatments T₆ (125 ppm) and T₁ (control) were observed at par with each other in number of branches.

At 65 DAS significant rise in number of branches was observed in treatment T₅ (100 ppm chitosan) tracked by treatment T₄ (75 ppm chitosan) when compared with treatment T₁ (control) and rest of the treatments under study. Treatments T₃ and T₂ remained at par with each other but found significantly superior when compared with control. Whereas, treatments T₆ (125 ppm) and T₁ (control) were found at par with each other.

At 85 DAS significantly superior number of branches were perceived in treatment T₅ (100 ppm) when compared with treatment T₁ (control) and rest of the treatments under study. Also treatments T₄ (75 ppm chitosan) and T₃ (50 ppm chitosan) followed superiority in case of number of branches when compared with treatment T₁ (control) and rest of the treatments. However, treatments T₂ (25 ppm chitosan) and T₆ (125 ppm chitosan) were found at par with each other but also found superior over treatment T₁ (control) in number of branches.

At harvest, highest number of branches were recorded in treatment T₅ (100 ppm chitosan) followed by treatments T₄ (75 ppm chitosan) and T₃ (50 ppm chitosan) when compared with treatment T₁ (control) and rest of the treatments under study. Treatment T₂ (25 ppm chitosan) estimated significant number of branches when correlated with treatment T₁ and left over treatment T₆, whereas, treatments T₆ and T₁ were found at par with each other.

Chitosan increased growth character viz., number of branches might be due to increased number of internodes or length of internodes because of increased cell number (Hong Yan and Shu Yu, 2001). This might be the reason for increase in number of branches in present investigation.

Meshram *et al.* (2018) applied varying concentrations of chitosan and IBA (25, 50, 75, 100 and 125 ppm) and concluded that application of 25 ppm of chitosan remarkably increased number of branches in soybean crop.

Leaf area plant⁻¹

Leaf area depends upon the number and size of leaves. Leaves play an important role in the absorption of light radiations and using it in photosynthetic process. Leaf size is influenced by light, moisture and nutrients. Hence, yield depends on leaf area of crop.

At 45 DAS leaf area plant⁻¹ was significantly influenced by different treatments. The range of leaf area recorded was 6.49-9.60 dm². At this stage significantly maximum leaf area plant⁻¹ was noted in treatment T₅ (100 ppm chitosan) followed by treatment T₄ (75 ppm chitosan) when compared with treatment T₁ (control) and other treatments. Also treatment T₃ (50 ppm chitosan) was found significantly superior over control T₁ (control). Whereas, treatment T₂ (25 ppm chitosan), T₆ (125 ppm chitosan) and T₁ (control) were found at par with each other in leaf area.

At 65 DAS the range of leaf area recorded was 11.39-16.03 dm² where treatment T₅ (100 ppm chitosan)

showed significantly highest leaf area plant⁻¹ followed by treatment T₄ (75 ppm chitosan) and T₃ (50 ppm chitosan) when compared with treatment T₁ (control) and rest of the treatments. While, treatment T₂ (25 ppm chitosan), treatment T₆ (125 ppm chitosan) and treatment T₁ (control) were found at par with each other in leaf area.

At 85 DAS significantly highest leaf area was witnessed in treatment T₅ (100 ppm chitosan) followed by treatment T₄ (75 ppm chitosan) when compared with treatment T₁ (control) and rest of the treatments. Treatment T₃ (50 ppm chitosan) also found superiorly significant when compared with treatment T₁ (control) and rest of the treatments. While treatment T₂ (25 ppm chitosan), treatment T₆ (125 ppm chitosan) and treatment T₁ (control) were found at par with each other in leaf area.

Chibu and Shibayama (2001) witnessed existence of greater chlorophyll content in the plant applied with chitosan. This contributed into the increase of the photosynthesis production and leaf area. The application of chitosan elevated key enzymes activities of nitrogen metabolism and improved transportation of nitrogen in functional leaves which might be the reason for enhanced leaf area in the present study.

Baraskar *et al.* (2018) tested four different plant growth regulators. Study was carried out to evaluate the impact of foliar sprays of plant growth regulators GA₃ (100,150,200 ppm), NAA (100,150,200 ppm), salicylic acid (500,1000,1500 ppm) and chitosan (100,125,150 ppm) on okra at three different concentrations and one control (without spray). Outcomes were application of 100 ppm chitosan impressively and significantly elevated leaf area when compared with control.

Total dry weight plant⁻¹

At 45 DAS the results of dry matter production specified that treatment T₅ (100 ppm chitosan) was found significantly superior followed by treatment T₄ (75 ppm chitosan) among all the treatments when compared with control and rest of the treatments under study. Treatments T₃ (50 ppm chitosan) and T₂ (25 ppm chitosan) had established significant results against treatment T₆ (125 ppm chitosan) and treatment T₁ (control). Whereas, treatments T₆ (125 ppm chitosan) also produced significantly higher dry matter over treatment T₁ (control).

At 65 DAS production of dry matter was significantly maximum in treatment T₅ (100 ppm chitosan) followed by treatments T₄ (75 ppm chitosan) and T₃ (50 ppm chitosan) when compared with treatments T₁ (control) and rest of the treatments under study. Correspondingly treatment T₂ (25 ppm chitosan) ensued significantly extra dry matter production when compared with treatment T₁ (control) and remaining treatments. Whereas treatment T₆ (125 ppm chitosan) was at par with treatment T₁ (control).

At 85 DAS dry matter was produced significantly maximum in treatment T₅ (100 ppm chitosan) followed by treatment T₄ (75 ppm chitosan) when compared with treatment T₁ (control) and rest of the treatments under study. Likewise treatments T₃ (50 ppm chitosan) and T₂ (25 ppm chitosan) supervised significantly extra dry matter production when

compared with treatment T₁ (control) and left over treatments. Whereas, treatment T₆ (125 ppm chitosan) stood at par with T₁ (control) in dry matter production.

Stimulating effect of chitosan on plant growth viz., dry matter may be attributed to an increase in the availability and uptake of water and essential nutrients through adjusting osmotic pressure and reducing the accumulation of harmful free radicals by increasing antioxidants and enzyme activities (Guan *et al.*, 2009). Higher area of leaves and chlorophyll content has contributed into the increase of the photosynthesis production which reflects significant amount of dry weight (Chibu and Shibayama, 2001). These might be the reasons for increase in dry matter in the present investigation.

Meshram *et al.* (2018) studied the efficacy of foliar application of chitosan and Indole-3-butyric acid on growth and productivity of soybean and stated that the application of chitosan @ 25 ppm improved dry weight in soybean crop significantly.

Relative growth rate (RGR)

Considering all the treatments under study, significantly maximum RGR was noted in treatment T₅ (100 ppm chitosan) i.e. 0.1017 g g⁻¹ day⁻¹ while, it was noted lowest in T₁ (control) i.e. 0.0556 g g⁻¹ day⁻¹ at 45-25 DAS. Treatment T₅ (100 ppm chitosan) earned highest RGR followed by treatments T₄ (75 ppm chitosan), T₃ (50 ppm chitosan) and T₂ (25 ppm chitosan) when compared with control and further treatments. Whereas, treatment T₆ (125 ppm chitosan) also gave significantly superior RGR when compared with treatment T₁ (control).

During 2nd phase i.e. 65-45 DAS treatment T₅ (100 ppm chitosan) exhibited significantly highest RGR when compared with treatment T₁ (control) and treatments left over. Correspondingly treatment T₄ (75 ppm chitosan) had also significantly higher RGR when compared with treatment T₁ (control) and remaining treatments under evaluation. However, treatments T₃ (50 ppm chitosan), T₂ (25 ppm chitosan), T₆ (125 ppm chitosan) were found at par with treatment T₁ (control) in RGR.

RGR for the duration of 85-65 DAS revealed that treatment T₅ (100 ppm chitosan) brought highest RGR when compared with control and remaining treatments under study. Treatments T₄ (75 ppm chitosan), T₃ (50 ppm chitosan), T₂ (25 ppm chitosan) and T₆ were found at par with treatment T₁ (control) in RGR.

Mondal *et al.* (2012) accomplished a field experiment to study the effect of different concentrations of chitosan (0, 50, 75, 125 and 150 ppm) on okra. Application of chitosan expressively increased relative growth rate by the application of 100 ppm chitosan.

Meshram *et al.* (2018) tested varying concentrations of chitosan and IBA (25, 50, 75, 100 and 125 ppm) and concluded that application of 25 ppm of chitosan remarkably enhanced RGR in soybean crop.

Net assimilation rate (NAR)

Net assimilation rate (NAR), synonymously called as unit leaf rate expresses the rate of dry weight increase at

Table1. Effect of chitosan on plant height, number of branches plant⁻¹ and leaf area plant⁻¹ in groundnut

Treatments	Plant height (cm)			Number of branches plant ⁻¹					Leaf area plant ⁻¹ (dm ²)				
	25 DAS	45 DAS	85 DAS	25 DAS	45 DAS	65 DAS	85 DAS	At harvest	25 DAS	45 DAS	85 DAS		
T ₁ (control)	10.68	26.34	38.78	40.74	1.30	1.90	2.50	3.35	3.80	1.79	6.49	11.39	12.11
T ₂ (25 ppm chitosan)	10.50	27.02	42.36	51.35	1.41	2.50	3.50	4.00	4.62	1.74	6.95	12.76	13.17
T ₃ (50 ppm chitosan)	10.33	27.16	42.91	52.39	1.58	2.50	3.50	4.50	4.64	1.73	7.03	13.61	15.01
T ₄ (75 ppm chitosan)	10.20	28.57	43.01	54.70	1.41	2.95	4.00	4.64	4.88	1.56	8.91	13.68	15.56
T ₅ (100 ppm chitosan)	10.38	30.82	46.15	57.95	1.60	3.0	4.30	5.00	5.00	1.61	9.60	16.03	16.94
T ₆ (125 ppm chitosan)	10.73	26.82	39.89	50.30	1.70	2.15	3.25	3.80	3.95	1.61	6.95	11.74	12.98
SE(m)±	0.460	0.851	0.625	0.564	0.097	0.044	0.100	0.112	0.173	0.053	0.322	0.512	0.524
CD at 5 %	—	2.548	1.866	1.683	—	0.117	0.294	0.329	0.503	—	0.958	1.527	1.560

Table2. Effect of chitosan on total dry weight plant⁻¹, RGR, NAR and yield plot⁻¹ in groundnut

Treatments	Total dry weight plant ⁻¹ (g)			RGR (g g ⁻¹ day ⁻¹)			NAR (g dm ⁻² day ⁻¹)			Yield plot ⁻¹ (kg)	
	25 DAS	45 DAS	85 DAS	45-25 DAS	65-45 DAS	85-65 DAS	45-25 DAS	65-45 DAS	85-65 DAS		
T ₁ (control)	2.58	7.85	29.96	39.01	0.0556	0.0355	0.0132	0.0671	0.0786	0.0382	0.550
T ₂ (25 ppm chitosan)	2.23	14.81	33.11	44.05	0.0947	0.0402	0.0147	0.1185	0.0956	0.0425	0.671
T ₃ (50 ppm chitosan)	2.19	15.00	34.08	46.25	0.0950	0.0410	0.0153	0.1694	0.0957	0.0442	0.736
T ₄ (75 ppm chitosan)	2.55	17.06	34.73	48.02	0.0962	0.0554	0.0162	0.1769	0.1141	0.0455	0.749
T ₅ (100 ppm chitosan)	2.38	18.20	37.93	49.27	0.1017	0.0670	0.0171	0.1799	0.1229	0.0487	0.831
T ₆ (125 ppm chitosan)	2.10	10.40	31.50	42.26	0.0800	0.0367	0.0143	0.0692	0.0794	0.0423	0.619
SE(m)±	0.064	0.398	1.289	1.00	0.0048	0.0034	0.0010	0.0040	0.0059	0.0012	0.0034
CD at 5 %	—	1.186	3.861	3.00	0.0119	0.0102	0.0030	0.0120	0.0176	0.0031	0.0101

any instant on a leaf area basis with leaf representing an estimate of the size of the assimilatory surface area (Gregory, 1926). Increase in NAR during reproductive phases might be due to increase efficiency of leaves for photosynthesis as a response of photosynthetic apparatus to increase demands for assimilates by growing seed fraction and also due to photosynthetic contribution by pod and sink demand on photosynthetic rate of leaves.

NAR at 45-25 DAS was significantly enhanced by treatments T₅ (100 ppm chitosan), T₄ (75 ppm chitosan) and T₃ (50 ppm chitosan) when compared with treatment T₁ (control) and remaining treatments under study. After that, treatment T₂ (25 ppm chitosan) also showed significantly higher NAR when compared with treatment T₆ (125 ppm chitosan). Where treatment T₆ (125 ppm chitosan) found at par with treatment T₁ (control) in NAR.

At 65-45 DAS significantly superior NAR was recorded in treatment T₅ (100 ppm chitosan) followed by T₄ (75 ppm chitosan) when compared with treatment T₁ (control) and treatments under examination. Treatments T₃ (50 ppm chitosan), T₂ (25 ppm chitosan), T₆ (125 ppm chitosan) were found at par with treatment T₁ (control) in NAR.

At 85-65 DAS NAR was found significantly maximum in treatment T₅ (100 ppm chitosan) followed by treatment T₄ (75 ppm chitosan), T₃ (50 ppm chitosan) and T₂ (25 ppm chitosan) when compared with treatment T₁ (control) and treatments under study. Whereas, treatment T₆ (125 ppm chitosan) also found significantly superior over treatment T₁ (control) in NAR.

Meshram *et al.* (2018) tried varying concentrations of chitosan and IBA (25, 50, 75, 100 and 125 ppm) and concluded that application of 25 ppm of chitosan remarkably enhanced NAR in soybean crop.

Shaheen *et al.* (2019) examined the effect of bio stimulants on plant growth and yield of potato. Foliar application of some plant growth stimulants was done (amino acid, 2.5 cm³ l⁻¹, chitosan 5 cm³ l⁻¹, potassium silicate 2 cm³ l⁻¹ and control treatment) for 3 times in 10 days interval starting at 40 days after planting date. The study revealed that application of chitosan @ 5 cm³ l⁻¹ significantly raised net assimilation rate.

Yield

Seed yield is influenced by morpho-physiological parameters such as plant height, leaf area and total dry matter production which are considered as yield contributing parameters.

Significantly maximum seed yield plot⁻¹ was produced in treatment T₅ (100 ppm chitosan) when compared with treatment T₁ (control) and rest of the treatments under study. Also treatments T₄ (75 ppm chitosan), T₃ (50 ppm chitosan), T₂ (25 ppm chitosan) and T₆ (125 ppm chitosan) were found significantly superior in a descending manner when compared with treatment T₁ (control) in production of seed yield plot⁻¹. The positive effect of chitosan on plant growth may be credited to an increase in the key enzyme activities of nitrogen metabolism and increased photosynthesis which enhanced plant growth (Gornik *et*

al., 2008; Mondal *et al.*, 2012). Chitosan applications significantly increased total carbohydrates accumulation in the fruits (Mawgoud *et al.*, 2010). These might be the reasons for increase in yield in the present study when compared with control and other treatments. The increase in groundnut yield due to chitosan application may be due to its effect in stimulating physiological processes, improving vegetative growth, followed by the active translocation of photo assimilates from source to sink tissue (Sharifa and Abu Muriefah, 2013). Chitosan when applied externally was observed to increase crop growth and ultimately yield.

REFERENCES

- Anonymous, 2018. Ann. report 2017-18 website www.krishi.maharashtra.gov.in.
- Baraskar, T. V., P. P. Gawande, N. V. Kayande, S.S. Lande and M. S. Naware, 2018. Effect of plant growth regulators on growth parameters of okra. *INTL J. Chem. studies.* 6(6): 165-168.
- Blackman, V. M. and J. Hamblin, 1919. Growth and development in physiology of crop plants, IInd Ed. Scientific publishers. Jodhpur. pp. 198-199.
- Chibu, H. and H. Shibayama, 2001. Effects of chitosan application on the growth of several crops. in: T. Uragami, K. Kurita and T. Fukamizo (Eds.), Chitin and chitosan in life science, Yamaguchi, Japan. pp. 235-239.
- Gorgery, F.G. 1926. The effect of climate conditions on growth of barley. *Ann. Bot.* 40 : 1-26
- Gornik, K., M. Grzesik and B.R. Duda, 2008. The effect of chitosan on rooting of grape vine cuttings and on subsequent plant growth under drought and temperature stress. *J. Fruit Ornament. Plant Res.* 16: 333-343.
- Guan, Y.J., J.Hu, X.J. Wang and C.X. Shao, 2009. Seed priming with chitosan improves maize germination and seedling growth in relation to physiological changes under low temperature stress. *J. Zhejiang University Sci B.* 10(6): 427-433.
- Hong Yan, W. and L. Shu Yu, 2001. The effect of chitosan and sodium alginate on the growth and photosynthesis of soybean. *J. North East Agric. Univ. China.* 8: 156-160.
- Mawgoud, A.M.R., A.S. Tantawy, M.A. El-Nemr and Y.N. Sassine, 2010. Growth and Yield Responses of Strawberry Plants to Chitosan Application. *European J. Sci. Res.* 39: 170-177.
- Meshram, S. D., R. D. Deotale, K. H. Chute, G. N. Jadhav and G. A. Padghan, 2018. Morpho-physiological and yield responses of soybean foliar sprays of chitosan and IBA. *J. Soils and Crops.* 28(1): 121-127.
- Mondal, M.M.A., M.A. Malek, A.B. Puteh, M.R. Ismail, M. Ashrafuzzaman and L. Naher, 2012. Effect of foliar application of chitosan on growth and yield in okra, *Australian J. Crop Sci.* 6(5): 918-921.
- Mondal, M. M. A., M.A. Malek, A.B. Puteh and M.R. Ismail, 2013. Foliar application of chitosan on growth and yield attributes of mungbean [*Vigna radiata* (L.) Wilczek]. *Bangladesh J. Bot.* 42(1): 179-183.
- Shaheen, A.M., M.E. Ragab, F.A. Rizk, H. Mahmoud, Soliman M.M. and N.M. Omar, 2019. Effect of some active stimulants on plant growth, tubers yield and nutritional values of potato plants grown in newly reclaimed soil. *J. Anim. and Plant Sci.* 29. 215-225.
- Sharifa, S. and Abu-Muriefah, 2013. Effect of chitosan on common bean (*Phaseolus vulgaris* L.) plants grown under water stress conditions. *INTL. Res. J. Agric. Sci. and Soil Sci.* 3(6): 192-199.
- Williams, R. S. 1946. The physiology of plant growth with special references to the concepts of net assimilation rate. *Ann. Bot.* 10 (10) : 41-42.